



6777 Nancy Ridge Drive
San Diego, CA 92121
Tel: 1.858.458.0866
Fax: 1.858.458.0833
Email: sales@genwaybio.com

Data Sheet

Fluorogenic DPP8 Assay Kit

Catalog #: GWB-53E1CE

DESCRIPTION: Dipeptidyl peptidase-8 (DPP8) is a ubiquitously expressed serine exopeptidase that cleaves X-proline dipeptides from the N-terminus of polypeptides. DPP8 shares homology and substrate recognition patterns with DPP4 and like DPP4, is thought to play a role in T-cell activation and immune function. The *Fluorogenic DPP8 Assay Kit* is designed to measure DPP8 activity using purified DPP8 for screening and profiling applications. It comes in a convenient 96-well format, with purified DPP8 enzyme, DPP substrate, and DPP assay buffer for 100 enzyme reactions. The key to the *Fluorogenic DPP8 Assay Kit* is the fluorogenic substrate. Using this kit, only one simple step on a microtiter plate is required for DPP8 reactions. The fluorometric substrate is incubated with a sample containing DPP8 enzyme to produce a fluorophore that can then be measured using a fluorescence reader.

COMPONENTS:

Ref #	Component	Amount	Storage	
80080	DPP8 human recombinant enzyme	5 µg	-80°C	Avoid freeze/ thaw cycles!
80300	DPP assay buffer	10 ml	-20°C	
80305	Fluorogenic DPP substrate 1 in DMSO (0.5 mM)	100 µl	-80°C	
	AMC Fluorescent standard (50 µM)	500 µl	-20°C	
	black, low binding NUNC black microtiter plate	1 plate	Room temp.	

APPLICATIONS: Great for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.458.0866** Fax **1.858.458.0833**
Or you can Email us at: sales@genwaybio.com
Please visit our website at: www.genwaybio.com



6777 Nancy Ridge Drive
San Diego, CA 92121
Tel: 1.858.458.0866
Fax: 1.858.458.0833
Email: sales@genwaybio.com

ASSAY PROTOCOL:

Immediately prior to assay:

- 1) Dilute DPP substrate 1 0.5 mM stock 5-fold with DPP assay buffer to make a 100 μ M solution. (Make only sufficient quantity needed for the assay; store remaining 0.5 mM stock solution in aliquots at -20°C.)
- 2) Dilute DPP8 in DPP assay buffer to 2 ng/ μ l (20 ng/reaction)*. Aliquot any remaining enzyme and store undiluted at -80°C. Keep diluted enzyme on ice. Discard any remaining diluted enzyme after use. *Note: optimal enzyme concentration may vary with the specific activity of the enzyme.
- 3) Dilute 25 μ l of the Fluorescent AMC standard (50 μ M stock) 2-fold with DPP assay buffer to make a 25 μ M solution. Make serial 2-fold dilutions of the fluorescent AMC standard in DPP buffer as follows: 12.5 μ M, 6.25 μ M, 3.12 μ M, 1.56 μ M, 0.78 μ M, 0.39 μ M, 0.19 μ M, 0.10 μ M. Aliquot the remaining 50 μ M AMC standard and store undiluted at -20°C.

Step 1:

In duplicate, add the reaction mixtures (below) to the microtiter black plate. Incubate at 22 °C for 10 min.

	Enzyme Positive Control	Test Inhibitor	AMC Standard Curve	Inhibitor Negative Control	"Blank" Negative Control
DPP8 (2 ng/ μ l)	10 μ l	10 μ l	-	-	-
DPP substrate 1 (100 μ M)	5 μ l	5 μ l	-	5 μ l	-
AMC standard (0.1 μ M – 50 μ M)	-	-	5 μ l	-	-
Inhibitor (in DPP assay buffer)	-	X μ l	-	X μ l	-
DPP assay buffer	85 μ l	85 - X μ l	95 μ l	95 - X μ l	100 μ l
Total	100 μl	100 μl	100 μl	100 μl	100 μl

Step 2:

Read sample in a microtiter-plate fluorimeter that is capable of excitation at wavelengths ranging from 350-380 nm and detection of emitted light ranging from 440-460 nm.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.458.0866** Fax **1.858.458.0833**


Or you can Email us at: sales@genwaybio.com

Please visit our website at: www.genwaybio.com



6777 Nancy Ridge Drive
San Diego, CA 92121
Tel: 1.858.458.0866
Fax: 1.858.458.0833
Email: sales@genwaybio.com

Example of Assay Results:



DPP8 enzyme activity, measured using the *Fluorogenic DPP8 Assay Kit*, GenWay Biotech, Cat No. *GWB-53E1CE*. *Note: Data shown is lot-specific. For lot-specific information, please contact GenWay Biotech at sales@genwaybio.com.*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.458.0866** Fax **1.858.458.0833**
Or you can Email us at: sales@genwaybio.com
Please visit our website at: www.genwaybio.com